



Case Report

The association of hyperthyroidism and immune thrombocytopenia: Are we still missing something?

Kalyan Mansukhbhai Shekhda*

Department of Medicine,
Glangwili General hospital (NHS
Wales), Carmarthen, Wales, UK

Received : 23-Sep-2017
Revised : 19-Dec-2017
Accepted : 14-Jan-2018

ABSTRACT

Though the American society of hematology Guidelines and British guidelines do not recommend screening of thyroid diseases in cases of immune thrombocytopenia (ITP), more than 160 cases of hyperthyroidism associated with ITP have been reported. Numerous case reports would suggest that patients with ITP and concurrent hyperthyroidism would respond to control of thyroid disease rather than the standard ITP treatment. Although this issue is still debatable, we report a case of a young female with a previous diagnosis of hyperthyroidism which was not well controlled, had presented with severe thrombocytopenia. Initially, all work-up had been done to find out the cause of thrombocytopenia. After all normal reports except deranged thyroid function tests, the patient was labeled as ITP and started on steroids. Even after a few months of steroid treatment, platelet counts had not improved. However, after starting antithyroid drugs, platelet counts had become normalized.

KEYWORDS: *Antithyroid drugs, Autoimmune disease, Platelet counts, Steroids*

INTRODUCTION

Though guidelines do not recommend the screening of thyroid diseases in cases of Immune thrombocytopenia (ITP), numerous case reports would suggest that patients with ITP and concurrent hyperthyroidism, would respond to control of thyroid disease rather than the standard ITP treatment.

CASE REPORT

We report a case of ITP in a known case of hyperthyroidism in a 36-year-old female of Indian origin. She initially presented to the outpatient department with asymptomatic thrombocytopenia.

However, she had a history of bleeding gums, menorrhagia, and ecchymotic patches on all over body 4 months back. These complaints were associated with weakness and easy fatigability for which she had consulted a private physician. Laboratory reports were suggestive of serum thyroid-stimulating hormone (TSH): 0.001 microIU/mL (normal range: 0.35–5.5 microIU/mL), hemoglobin: 4.8 g%, white blood cells 5000/ μ L, and platelet: 36,000/ μ L. Considering anemia due to an acute bleeding episode, she was given 2-pack cell volume. After a week, she underwent laboratory investigations in which thrombocytopenia was persistent. She consulted a hematologist for that report and underwent investigations for thrombocytopenia. All investigations turned out to be negative (serum lactate dehydrogenase, direct Coombs test (DCT) and indirect DCT = Negative.) Even though reports were suggestive of hyperthyroidism, antithyroid

medications (carbimazole 10 mg once a day) were stopped. She was started on steroids; however, there was no improvement in platelet counts. After 3 months of unsatisfactory treatment, she presented at our center. On general examination, the patient was conscious, cooperative, and well oriented to time, place, and person. Pulse rate was 110/min with the blood pressure of 120/70 mmHg. There was no edema, icterus, cyanosis, clubbing, and lymphadenopathy. On thyroid gland examination, there was no thyroid enlargement and bruits present. Per abdomen examination spleen and liver were not palpable and the rest of systemic examinations were unremarkable.

On admission, laboratory investigations revealed: hemoglobin 13.6 g/dL, total leukocyte count 7800/ μ L, platelet count 21,000/ μ L. Serum TSH 0.01 mIU/L (normal range: 0.35–5.5 mIU/L), serum free T3 8.48 pmol/L (normal range: 3.542–6.468 pmol/L), and serum free T4 37.97 pmol/L (normal range 11.5–22.7 pmol/L). Urine routine examination, liver function test, and renal function test were within normal limits. Peripheral smear did not show malarial parasite. Dengue non-specific antigen 1, Ig-G, and Ig-M were negative. Human Immunodeficiency Virus (HIV), HbsAg, hepatitis C virus were

*Address for correspondence:

Dr. Kalyan Mansukhbhai Shekhda,
Department of General Medicine, Glangwili General Hospital (NHS Wales),
Carmarthen, SA31 2AF, Wales, UK.
E-mail: kalokly@gmail.com

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Shekhda KM. The association of hyperthyroidism and immune thrombocytopenia: Are we still missing something? Tzu Chi Med J 2018;30:188-90.

Access this article online	
Quick Response Code: 	Website: www.tcmjmed.com
	DOI: 10.4103/tcmj.tcmj_139_17

negative Antinuclear antibody profile was negative ruling out autoimmune causes. Bone marrow examination revealed normal functioning marrow. However, we restarted carbimazole 10 mg three times a day which had been kept on hold, and steroids were tapered off. After starting antithyroid medications, her platelet counts gradually improved. During subsequent follow-up, her thyroid function tests and platelet counts became normal. The trend of her platelet counts and thyroid function tests is summarized in [Tables 1 and 2].

DISCUSSION

Coexistence of ITP and hyperthyroidism is a well-documented but not a well-understood association. The association of these two diseases can be explained by the theory of the production of two kinds of antibodies (platelet-associated IgG, antithyroid antibodies [Microsomal and thyroglobulin antibodies]) against platelets and TSH receptor [1]. In addition to this, shorter platelet lifespan, which is metabolic in nature rather than immunogenic, in patients with hyperthyroidism and increased reticuloendothelial phagocytic activity are also postulated as a mechanism for thrombocytopenia in a case of hyperthyroidism [2,3]. The presentation of this association varies from case to case. It can present with a new onset of thrombocytopenia without life-threatening symptoms in a patient with preexisting hyperthyroidism, or it can present with a life-threatening thyrotoxic crisis which mimics sepsis in a known case of stable ITP [4,5]. In fact, studies show that autoimmune thrombocytopenia, in the course of thyroid disease, is far more prevalent than vice versa [1].

Although hyperthyroidism is known to shorten the platelet survival by enhancing reticuloendothelial activity, the resultant thrombocytopenia is usually mild-to-moderate [3]. Therefore,

this mechanism cannot explain the severe thrombocytopenia of the present case. The severity of the thrombocytopenia and its responsiveness to antithyroid medication strongly support the first view that both disorders stemmed from a common autoimmunity. A similar response was seen in a case of ITP with Graves, whereby dose adjustment of methimazole normalized the thyroid function as well as spontaneous recovery of platelet count which had fallen due to irregular adherence to antithyroid medication [4]. However, in our case-patient was already prescribed steroids for ITP for 2 months, but the failure of the standard therapy to improve platelet counts prompted us to look for other therapeutic options. After correcting thyroid function with antithyroid medication, platelet count raised without concurrent steroid supplementation. This suggests that in a patient of hyperthyroidism with ITP, correction of the hyperthyroidism plays a role in correcting the immune system imbalance which impairs not only the thyroid function but also the platelet antigen response. Despite more than 160 cases of hyperthyroidism associated with ITP that have been reported, the American society of hematology Guidelines and British guidelines do not recommend screening of thyroid diseases in cases of ITP [6-8]. This is might be because of lack of clarity of the exact mechanism and association.

Declaration of patient consent

The author certifies that the patient has obtained appropriate patient consent form. In the form, the patient has given her consent for her images and other clinical information to be reported in the journal. The patient understands that her name and initial will not be published and due efforts will be made to conceal her identity, but anonymity cannot be guaranteed.

Table 1: Trend of thyroid function test and platelet counts before starting antithyroid therapy

Date	Serum TSH (normal range: 0.35-5.5 mIU/L)	Free T3 (normal range: 3.542-6.468 pmol/L)	Free T4 (normal range: 11.5-22.7 pmol/L)	Platelet counts (normal range: 150,000-450,000/ μ L)	Treatment received	Steroid dose
March 08, 2016	0.001	7.3	29.5	36,000	FFP, PCV, steroids	Methylprednisolone 1 g/day
March 16, 2016	0.06	8.12	30.1	12,000	Steroids	for 3 days then prednisolone
April 08, 2016	-	-	-	73,000	Steroids	70 mg/day (1 mg/kg/day) for
May 05, 2016	0.08	-	-	45,000	Steroids	1 st month then tapered. 5 mg
June 01, 2016	-	-	-	121,000	Steroids	reduced every week
July 08, 2016	0.05	12.2	35.3	34,000	Steroids	
July 14, 2016	0.01	8.48	37.97	21,000	Carbimazole	

TSH: Thyroid-stimulating hormone, T3: Triiodothyronine, T4: Thyroxine, FFP: Fresh frozen plasma, PCV: Pack cell volume

Table 2: Trend of thyroid function test and platelet counts after starting antithyroid therapy

Date	Serum TSH (normal range: 0.35-5.5 mIU/L)	Free T3 (normal range: 3.542-6.468 pmol/L)	Free T4 (normal range: 11.5-22.7 pmol/L)	Platelet counts (normal range: 150,000-450,000/ μ L)	Treatment received	Dose of carbimazole
July 17, 2016	-	-	-	30,000	Carbimazole	30 mg
August 08, 2016	0.02	3.90	13.53	292,000	Carbimazole	
September 19, 2016	0.13	4.32	14.33	308,000	Carbimazole	
October 25, 2016	0.6	5.01	18.25	210,000	Carbimazole	
December 05, 2016	0.7	5.4	20.72	249,000	Carbimazole	
January 06, 2017	-	4.3	17.85	297,000	Carbimazole	
February 13, 2017	-	-	22.45	271,000	Carbimazole	

TSH: Thyroid-stimulating hormone, T3: Triiodothyronine, T4: Thyroxine

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Cordiano I, Betterle C, Spadaccino CA, Soini B, Girolami A, Fabris F, et al. Autoimmune thrombocytopenia (AITP) and thyroid autoimmune disease (TAD): Overlapping syndromes? *Clin Exp Immunol* 1998;113:373-8.
2. Panzer S, Haubenstock A, Minar E. Platelets in hyperthyroidism: Studies on platelet counts, mean platelet volume, ¹¹¹-indium-labeled platelet kinetics, and platelet-associated immunoglobulins G and M. *J Clin Endocrinol Metab* 1990;70:491-6.
3. Kurata Y, Nishioeda Y, Tsubakio T, Kitani T. Thrombocytopenia in Graves' disease: Effect of T3 on platelet kinetics. *Acta Haematol* 1980;63:185-90.
4. Sugimoto K, Sasaki M, Isobe Y, Tamayose K, Hieda M, Oshimi K, et al. Improvement of idiopathic thrombocytopenic purpura by antithyroid therapy. *Eur J Haematol* 2005;74:73-4.
5. Azar M, Frates A, Rajput V. Idiopathic thrombocytopenic purpura (ITP) and hyperthyroidism: An unusual but critical association for clinicians. *J Hosp Med* 2008;3:431-3.
6. Aggarwal A, Doolittle G. Autoimmune thrombocytopenic purpura associated with hyperthyroidism in a single individual. *South Med J* 1997;90:933-6.
7. George JN, Woolf SH, Raskob GE, Wasser JS, Aledort LM, Ballem PJ, et al. Idiopathic thrombocytopenic purpura: A practice guideline developed by explicit methods for the American Society of Hematology. *Blood* 1996;88:3-40.
8. British Committee for Standards in Haematology General Haematology Task Force. Guidelines for the investigation and management of idiopathic thrombocytopenic purpura in adults, children and in pregnancy. *Br J Haematol* 2003;120:574-96.

