



Case Report

Pelvic myoma arising from vaginal cuff after hysterectomy: A case report and literature review

Yi-Jen Ho^a, Pei-Chen Li^a, Chiu-Hsuan Cheng^b, Dah-Ching Ding^{a,c*}

^aDepartment of Obstetrics and Gynecology, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation and Tzu Chi University, Hualien, Taiwan,

^bDepartment of Pathology, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation and Tzu Chi University, Hualien, Taiwan,

^cInstitute of Medical Sciences, Tzu Chi University, Hualien, Taiwan

Submission : 05-Aug-2019

Revision : 15-Sep-2019

Acceptance : 03-Oct-2019

Web Publication : 05-Dec-2019

ABSTRACT

Leiomyomas are the most common benign gynecologic tumors and the most common surgical indication for hysterectomy or myomectomy. Recurrent pelvic leiomyomas or vaginal leiomyomas are rare. We report the case of a 60-year-old woman with a history of a cesarean total hysterectomy 25 years before presenting to us, who developed a pelvic leiomyoma arising from the vaginal cuff. The patient underwent single-port laparoscopic tumor excision and recovered uneventfully.

KEYWORDS: Cuff, Hysterectomy, Laparoscopy, Myoma, Recurrence

INTRODUCTION

Uterine leiomyomas are the most common benign gynecologic tumors and the most common surgical indication for hysterectomy or myomectomy. However, recurrent pelvic myomas after morcellation during laparoscopy have been reported at rates of 0.12%–0.95% [1]. Furthermore, low-incidence leiomyosarcomas can occur under this setting [2]. Thus, in-bag manual morcellations have been recommended [3].

Recurrent myomas after a total hysterectomy are rare with <110 cases reported until now [4]. Here, we report the case of a woman with a pelvic recurrent myoma arising from the vaginal cuff after total hysterectomy during a cesarean section that had taken place 25 years before.

CASE REPORT

A 60-year-old woman, gravida 3, para 3, presented with an asymptomatic pelvic mass. She had undergone cesarean total hysterectomy with bilateral adnexa preservation 25 years before due to the presence of a uterine myoma. After that, she experienced surgical menopause. Her medical history included coronary artery disease, type 2 diabetes mellitus, dyslipidemia, and hyperparathyroidism. She had no history of hormone replacement therapy after menopause.

She was referred to our hospital with a pelvic mass found incidentally during a routine health examination. A whole-body magnetic resonance (MR) image showed a well-circumscribed mass of 7.2 × 6.4 cm in the pelvis [Figure 1a and

b]. Vaginal examination revealed an upward deviation of her vaginal cuff. Transvaginal ultrasound images demonstrated a 6.25 × 5.03 × 6.94 cm pelvic tumor [Figure 1c and d]. Tumor markers (CA 125, CA 19-9, and CEA) were all within normal ranges.

We scheduled the patient for surgery to rule out the possibility of malignancy. She underwent a single-port laparoscopic tumor excision. During the surgery, we found the tumor to be pedunculated and originating from the vaginal cuff [Figure 2a]. The tumor was round, firm, and sharply circumscribed resembling a uterine leiomyoma [Figure 2b]. A frozen section revealed a leiomyoma [Figure 3a and b]. The immunohistochemistry was positive for vimentin [Figure 3c] and smooth muscle actin (SMA) [Figure 3d], which are typical of myomas. We discharged the patient 2 days after the operation. Her condition was good when she came back to our outpatient department for a follow-up 1 month after surgery.

DISCUSSION

Uterine leiomyomas are the most common pelvic tumors in women. They are benign and monoclonal tumors originating from uterine smooth muscle cells. Interestingly, they can behave

**Address for correspondence:*

Dr. Dah-Ching Ding,
Department of Obstetrics and Gynecology, Hualien Tzu Chi Hospital,
Buddhist Tzu Chi Medical Foundation, 707, Section 3, Chung-Yang Road,
Hualien, Taiwan.
E-mail: dah1003@yahoo.com.tw,

Access this article online

Quick Response Code:



Website: www.tcmjmed.com

DOI: 10.4103/tcmj.tcmj_169_19

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Ho YJ, Li PC, Cheng CH, Ding DC. Pelvic myoma arising from vaginal cuff after hysterectomy: A case report and literature review. Tzu Chi Med J 2020; 32(4): 398–400.




Figure 1: Image study of the pelvic myoma. Pelvic magnetic resonance image showing sagittal (a) and coronal (b) views of the myoma. Pelvic ultrasound showing sagittal (c) and coronal (d) views of the myoma




Figure 2: Laparoendoscopic surgery for pelvic myoma. (a) Pelvic myoma along the vaginal cuff region. (b) Myoma retrieval by in-bag manual morcellation




Figure 3: Pathology of pelvic myoma. (a and b) Histology of the myoma showed by H and E staining. Smooth muscle cells were elongated with eosinophilic cytoplasm. Smooth muscle bundle showed fascicular pattern. The immunohistochemistry was positive for vimentin (c) and smooth muscle actin (d) which are typical of myoma. Scale bar = 10 μ m (b) and = 20 μ m (a, c, and d)

like malignancies and metastasize to various sites outside the uterus, such as the pelvis, lungs, retroperitoneum, heart, vessels, spine, and lymph nodes [5]. Vaginal leiomyomas are rare with about 300 reported cases [6]. A vaginal vault leiomyoma developing after a hysterectomy is even more uncommon, and only six cases have been reported [7-12]. The recurrence

intervals have ranged from 1 to 25 years [Table 1]. In our case, the myoma recurred 25 years after the hysterectomy.

Vaginal leiomyomas are pathologically similar to uterine leiomyomas. They are thought to arise from the vascular smooth muscle of the vagina, the submucosal smooth muscle, the smooth muscle of the round ligament, or from mesenchymal stem cell differentiation into smooth muscle cells [13]. Peritoneal seeding after myomectomy or hysterectomy during laparotomy or laparoscopic morcellation for uterine leiomyomas is another possible cause of these tumors. In theory, the residual morcellation fragments may be introduced to the peritoneum or adjacent tissues, where they later develop into pelvic or vaginal leiomyomas [5]. The growth of vaginal leiomyomas may depend on the association between hormones and locally active growth factors. They show strong positivity for SMA. In the study by Zhao *et al.*, 38.5% and 34.6% of vaginal leiomyomas expressed estrogen receptor and epidermal growth factor receptor, respectively, suggesting a role for estrogen in the pathogenesis of the tumor. However, recurrent leiomyomas can occur in women with poor estrogenic statuses even after a total hysterectomy and bilateral salpingo-oophorectomy without hormone replacement therapy [10].

Vaginal leiomyomas may be asymptomatic. However, symptoms may be present depending on the site of occurrence and include lower abdominal pain, low back pain, vaginal bleeding, dyspareunia, urinary frequency dysuria, and urinary obstruction [6].

The diagnosis of vaginal leiomyomas through ultrasonography may be difficult, and MR imaging is a superior tool for diagnosing them. In MR imaging, these tumors can appear as single, smooth tumors with homogeneous low-signal intensity on both T1- and T2-weighted images, as is typical in uterine leiomyomas. However, the cellular and vascular-type leiomyomas may show hyperintensity on T2-weighted images and marked contrast enhancement [14]. Malignant changes of primary leiomyomas or malignancies have also been reported [7]. Histopathological confirmation is the gold standard of diagnosis to rule out the possibility of malignancy.

Surgical resection of the tumor can be performed vaginally, laparoscopically, or abdominally depending on the tumor size, location, and accessibility. In our case, the leiomyoma originated from the vaginal cuff and protruded into the pelvic cavity. We resected it successfully through a laparoscopic approach. Vaginal myomectomy and transvaginal natural orifice transluminal endoscopic surgery myomectomy are feasible alternatives. Oophorectomy may be performed concurrently with the removal of the fibroids, ensuring an estrogen-poor environment and reducing the risk of recurrence. Complete and in-bag removal of the tumors with undetermined pathology is recommended during the operation to prevent inadvertent dissemination or spillage of occult malignancies.

CONCLUSION

Although rare, vaginal cuff leiomyomas can develop even in hysterectomized women. The outcome is favorable after complete excision of the tumor.

Table 1: Summary of cases with vaginal cuff leiomyomas after hysterectomy

Author (years)	Patient age (years)	Time to diagnosis (years)	Hysterectomy indication/procedure	Recurrence site	Tumor size (cm)	Pathology	Treatment
Yogesh <i>et al.</i> , 2005 [7]	42	5	Leiomyoma/TAH	Anterior vaginal wall	19×11×9	Leiomyoma	Surgical excision with BSO (laparotomy)
Yanamandra <i>et al.</i> , 2007 [8]	47	8	Leiomyoma/TAH	Vaginal vault	15×10	Leiomyoma	Surgical excision with BSO (laparotomy)
Ismail and Adams 2009 [9]	47	1	Leiomyoma/TAH	Vaginal vault	3×3	Leiomyoma	Surgical excision (vaginal)
Yarci <i>et al.</i> , 2010 [10]	70	25	Leiomyoma/TAH + BSO	Vaginal vault	7×3	Leiomyoma	Surgical excision (vaginal)
P <i>et al.</i> , 2014 [11]	48	6	Leiomyoma/TAH	Vaginal stump near right broad ligament	4×4	Leiomyoma	Surgical excision (laparotomy)
Bhuyar 2015 [12]	65	4	Leiomyoma/TAH + BSO	Vaginal vault	10×10	Atypical leiomyoma	Surgical excision (vaginal)
Present case	65	25	Leiomyoma/cesarean hysterectomy	Vaginal vault	7×6	Leiomyoma	Surgical excision (laparoscopic)

TAH: Total abdominal hysterectomy; BSO: Bilateral salpingo-oophorectomy

Declaration of patient consent

We, the authors, certify that we have obtained all appropriate patient consent forms. In the forms, the patient has given her consent for her images and other clinical information to be reported in the journal. The patient understands that her names and initials will not be published and due efforts will be made to conceal her identity, but that anonymity cannot be guaranteed.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Van der Meulen JF, Pijnenborg JM, Boomsma CM, Verberg MF, Geomini PM, Bongers MY. Parasitic myoma after laparoscopic morcellation: A systematic review of the literature. *BJOG* 2016;123:69-75.
2. Wright JD, Tergas AI, Cui R, Burke WM, Hou JY, Ananth CV, et al. Use of electric power morcellation and prevalence of underlying cancer in women who undergo myomectomy. *JAMA Oncol* 2015;1:69-77.
3. Ding DC, Hong MK, Chu TY, Chang YH, Liu HW. Robotic single-site supracervical hysterectomy with manual morcellation: Preliminary experience. *World J Clin Cases* 2017;5:172-7.
4. Kyriakopoulos K, Domali E, Stavrou S, Rodolakis A, Loutradis D, Drakakis P. Recurrent benign leiomyomas after total abdominal hysterectomy. Rich or poor estrogenic environment may lead to their recurrence? *Int J Surg Case Rep* 2018;44:191-3.
5. Barnaś E, Książek M, Raś R, Skręt A, Skręt-Magierło J, Dmoch-Gajzlarska E. Benign metastasizing leiomyoma: A review of current literature in respect to the time and type of previous gynecological surgery. *PLoS One* 2017;12:e0175875.
6. Chakrabarti I, De A, Pati S. Vaginal leiomyoma. *J Midlife Health* 2011;2:42-3.
7. Yogesh K, Amita M, Rajendra K, Raju A, Rekha W, Hemant T. Vaginal leiomyoma developing after hysterectomy - Case report and literature review. *Aust N Z J Obstet Gynaecol* 2005;45:96-7.
8. Yanamandra SR, Redman CW, Coomarasamy A, Varma R. Leiomyoma appearing in the vaginal vault following hysterectomy. *J Obstet Gynaecol* 2007;27:91-2.
9. Ismail SM, Adams SA. Vaginal vault leiomyoma. *J Obstet Gynaecol* 2009;29:160-1.
10. Yarci A, Bayramov V, Sükür YE, Yüce T, Berker B. Vaginal vault leiomyoma: 25 years after total abdominal hysterectomy. *J Minim Invasive Gynecol* 2010;17:116-7.
11. Pallavee P, Ghose S, Samal S, Begum J, Zabeen M. Fibroid after hysterectomy: A diagnostic dilemma. *J Clin Diagn Res* 2014;8:OD01-2.
12. Bhuyar S. Recurrent atypical leiomyoma arising from vaginal cuff after hysterectomy-A rare case report. *IOSR-J Dental Med Sci* 2015;14:69-72.
13. Zhao Y, Li Y, Xu Y. Clinico-pathologic analysis of 26 cases of leiomyoma of the vagina. *Beijing Da Xue Xue Bao Yi Xue Ban* 2003;35:37-40.
14. Shimada K, Ohashi I, Shibuya H, Tanabe F, Akashi T. MR imaging of an atypical vaginal leiomyoma. *AJR Am J Roentgenol* 2002;178:752-4.