



## Review Article

# Chinese herbs and acupuncture to improve cognitive function in Alzheimer's disease

Wei WuLi<sup>a,b</sup>, Horng-Jyh Harn<sup>a,c</sup>, Tzyy-Wen Chiou<sup>b</sup>, Shinn-Zong Lin<sup>a,d,\*</sup>

<sup>a</sup>Bioinnovation Center, Buddhist Tzu Chi Medical Foundation, Hualien, Taiwan, <sup>b</sup>Department of Life Science, National Dong Hwa University, Hualien, Taiwan, <sup>c</sup>Department of Pathology, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, Hualien, Taiwan, <sup>d</sup>Department of Neurosurgery, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, Hualien, Taiwan

**Submission** : 27-Mar-2020  
**Revision** : 07-May-2020  
**Acceptance** : 22-May-2020  
**Web Publication** : 06-Feb-2021

### ABSTRACT

Alzheimer's disease (AD) is a neurodegenerative disease with a variety of causes. Traditional Chinese medicine (TCM), which includes the two main approaches of acupuncture and herbal medication, views the human body as a self-controlled system network. Fundamental theories, including “qi,” the five elements, and the theory of viscera, form the basis for classification. Diseases in humans are considered to be caused by an imbalance of “yang qi” and “yin qi” that lead to the nonhomeostasis of organs. Acupuncture is derived from 12 main meridians and 365 acupuncture points characterized by “blood and qi.” Needling of different positions corresponds to specific disease treatments to increase qi. Treatment with Chinese herbal medicines is based on syndrome differentiation characterized as “Zheng” which differs from the cause orientation approach of Western medicine. In this article, we review basic and clinical research studies that describe TCM herbs and acupuncture for the treatment of AD. Moreover, we propose that these two approaches be integrated to improve the outcomes for AD patients.

**KEYWORDS:** *Acupuncture, Alzheimer's disease, Qi, Traditional Chinese medicine, Zheng*

## INTRODUCTION

Alzheimer's disease (AD) features complex pathogenesis involving multiple pathophysiological processes and has become a serious public health problem worldwide [1]. The complex mechanisms of disease and the continued failure of drug development have created an urgent need for additional treatment options for AD [2].

Traditional Chinese medicine (TCM) using herbal products have been proven to prevent and treat dementia [3,4], and many small molecules that effectively inhibit AD have been extracted from Chinese herbs [5]. Chinese herbal medicines show potential for the treatment of AD. Acupuncture is another standard therapeutic approach of TCM. Using the scientific method, there has been continued exploration of Chinese herbal medicines combined with acupuncture to increase the practical benefits of AD treatment.

### Beta-amyloid accumulation

The major hypotheses of AD include the amyloid hypothesis and the tau hypothesis. The genes associated with beta-amyloid deposits include amyloid-beta precursor protein (APP) [6] and β-secretase-1 (BACE-1) [7], while the genes that primarily regulate γ-secretase are presenilin-1 (PSEN-1) and presenilin-2 (PSEN-2) [8]. Under normal conditions, the APP is cleaved by α-secretase to produce a peptide of 83 amino acid residues that

is then further processed by γ-secretase to form the P3 peptide fragment (24 or 26 residues). Peptides of this length are easily metabolized. Deregulation of these genes can lead to enhanced abnormal cleavage and the formation of amyloid peptides that are not easily metabolized (40 or 42 residues) [9].

### Tau protein phosphorylation

Tau is a microtubule-associated protein that is abundant in the central nervous system. In neurodegenerative disease such as AD and Parkinson's disease, Tau aggregates are commonly found, forming neurofibrillary tangles in the brain [10].

### Other hypotheses

Among patients diagnosed with AD, only 25% develop dementia caused by limbic predominant age-related TDP-43 encephalopathy (LATE) [11]. Early-onset AD is usually attributed to genetic mutation. These genetic mutations include APP BACE-1, PSEN-1, and PSEN-2. However, 95% of AD patients have late-onset AD, which is not necessarily caused by genetic mutation [12]. The complex natures of this disorder and the continued failure to develop effective drugs for AD have created an urgent need for more treatment options.

*\*Address for correspondence:*

Dr. Shinn-Zong Lin,  
 Department of Neurosurgery, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, 707, Section 3, Chung-Yang Road, Hualien, Taiwan.  
 E-mail: shinnzong@yahoo.com.tw

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

**For reprints contact:** WKHLRPMedknow\_reprints@wolterskluwer.com

**How to cite this article:** WuLi W, Harn HJ, Chiou TW, Lin SZ. Chinese herbs and acupuncture to improve cognitive function in Alzheimer's disease. Tzu Chi Med J 2021; 33(2): 122-7.

Access this article online	
<b>Quick Response Code:</b> 	<b>Website:</b> www.tcmjmed.com
	<b>DOI:</b> 10.4103/tcmj.tcmj_51_20

## TRADITIONAL CHINESE MEDICINE HYPOTHESIS: ZHENG

In TCM, the syndrome known as “Zheng” is a major concept in the diagnosis of disease. Zheng can be understood through the following basic features: (1) Visual observation of the appearance of the tongue and detection of the pulse condition reflect disturbances of the “Zang-Fu” organs, channels and collaterals, and qi and blood in the body. (2) Different illnesses may have the same Zheng. Through the visual, auditory, and tactile senses the physician observes the patient’s appearance and asks questions regarding the patient’s feelings, life habits, and physiological changes. The major identification methods of Zheng comprise eight theoretical perspectives: Qi, blood, body fluids, zang-fu, channel collateral, six-channel, Wei-Qi-Ying-blood, sanjiao, and disease causes [13-16].

### Qi


Chinese medicine combines Chinese philosophy with rules of thumb and treats diseases by natural methods. In the formation and development of Chinese medicine, the concept of “qi,” often translated as vital energy, is the backbone of the entire system. This includes two main points. The first is the dynamic balance between people and the outside environment; for example, the different seasons and changes in the weather produce wind, cold, heat, moisture, dryness, and fire. When the immune system is weakened, people are unable to bear the natural changes and become sick. The second regards to health protection. Qi flows through the circulatory system, with a strengthening effect on the corresponding organs that are active at different time points. The time period from 1 AM to 3 AM, for example, is considered the most vigorous time for liver metabolism; if one does not rest during this time, the burden on the liver will increase. There are many methods to regulate the circulation of qi in the body, with the major methods being Chinese herbs and acupuncture. Indeed, the National Institutes of Health has acknowledged that acupuncture therapy is an effective treatment for several common diseases, such as back pain, knee pain, headaches, and osteoarthritis [17,18].

### Traditional Chinese medicine hypothesis for dementia

TCM explained as a cybernetic approach regards the human body as a self-controlled system network. TCM use qi, five elements, and the theory of viscera as a basis for classification. Chinese herbal medicine also features the same classification system. The theory of Chinese medicine views dementia as a result of the shrinking of the brain. TCM has put forward a theory that “kidneys give rise to the marrow, and the brain is the sea of marrow” [Figure 1]; thus, deficiency in kidney functions will lead to dementia [19-21]. Chinese herbs and acupuncture may increase the chances of treating AD. Continued exploration of Chinese herbal medicine and acupuncture treatment will help provide novel directions for the development of AD drugs and offering practical treatment benefits.

## EFFECTS AND MECHANISMS OF ACTION OF SMALL-MOLECULE DRUGS EXTRACTED FROM CHINESE HERBS IN AD TREATMENT

The earliest pharmacopoeia from China is the Shen Nong Ben Cao Jing (“Divine Farmer’s Pharmacopoeia”) [22]. In CA 1596, Li Shi-Zhen compiled the Ben Cao Gang Mu (“Pharmacopoeia



**Figure 1:** Five elements corresponding to different body organs. Chinese medicine classifies the five internal organs by five elements. Chinese herbal medicines are also classified using the same system. The brain is considered to be closely related to the function of the kidneys. Chinese medicine teaches that the bone marrow, kidneys, and brain function closely together. Kidney deficiency will lead to less bone marrow, which leads to insufficient blood and cerebral dysfunction and ultimately dementia

Classified by Section and Sub-section”). Li’s consolidation of the previous pharmacopoeia together with his own views and observations, created a more thorough pharmacopoeia [23]. TCM classifies Chinese herbs according to qi and five elements theory. The brain and kidneys are categorized in the water element. We summarize the effects and possible mechanisms of small-molecule drugs extracted from 10 commonly used TCMs and the effects of these molecules on meridians and organs in the AD setting.

### Epimedii folium (淫羊藿)

Epimedii Folium (Yin yang huo; 淫羊藿) is a major therapy for kidney yang deficiency syndrome, and it can improve androgen hormone levels as well as combat wind qi, damp qi, and cold qi blocking circulation [24,25].

### Radix polygalae (遠志)

Radix polygalae (“Yuan Zhi,” 遠志) is a medicinal plant mainly used for the treatment of “insomnia with syndrome of incoordination between heart and kidney” and to improve memory [26-28].

### Morinda officinalis (巴戟天)

Morinda officinalis (“Ba Ji Tian,” 巴戟天) is a TCM used to tonify the kidney and strengthen yang. Morinda officinalis has hypoglycemic and antioxidant activities [29].

### Acorus tatarinowii schott (石菖蒲)

Acorus tatarinowii Schott (“Shi Chang Pu,” 石菖蒲) has mainly been used for the treatment of neuron-associated and digestive diseases in the clinical practice of TCM [30]. It can also improve the viability of cardiac myocytes and contribute to therapeutic effects on dementia [31,32].

### Rehmannia glutinosa (地黃)

Rehmannia glutinosa (“Dihuang,” 地黃) can invigorate the kidney, “nourishing yin clearing heat,” and also ameliorates learning and memory impairment [33-35].

### Glycyrrhiza uralensis (甘草)

Glycyrrhiza uralensis (“Gan Cao,” 甘草), is an herbal medicine used for the treatment of sore throat, cough, bronchitis, peptic ulcers, arthritis, and allergic diseases [36].

**Table 1. The system of meridians and collaterals**

Meridians			
<b>□. Twelve meridians (十二經脈)</b>			
1.Three yin meridians of hand(手三陰經)	2.Three yang meridians of hand(手三陽經)	3.Three yin meridians of foot(足三陰經)	4.Three yang meridians of foot(足三陽經)
1.1. Tai-yin lung Meridian of hand(手太陰肺經)	2.1. Yang-ming large intestine meridian of hand(手陽明大腸經)	3.1. Tai-yin spleen meridian of foot(足太陰脾經)	4.1. Yang-ming stomach meridian of foot(足陽明胃經)
1.2. Shao-yin heart meridian of hand(手少陰心經)	2.2. Tai-yang small intestine meridian of hand(手太陽小腸經)	3.2. Shao-yin kidney meridian of foot(足少陰腎經)	4.2. Tai-yang bladder meridian of foot(足太陽膀胱經)
Jue-yin pericardium 1.3 meridian of hand(手厥陰心包經)	2.3. Shao-yang Sanjiao meridian of Hand(手少陽三焦經)	3.3. Jue-yin liver meridian of foot(足厥陰肝經)	4.3. Shao-yang gallbladder meridian of foot(足少陽膽經)
<b>□. Eight extra meridians (奇經八脈)</b>			
1.Conception vessel(任脈)	2.Governor vessel(督脈)	3.Chong Vessel(沖脈)	4.Belt vessel(帶脈)
5.Yin Link Vessel(陰維脈)	6.Yang Link Vessel(陽維脈)	7.Yin Heel Vessel(陰蹻脈)	8.Yang Heel vessel(陽蹻脈)
□. Twelve divergent channels (十二經別). □. Twelve sinew channels (十二經筋). □. Twelve cutaneous regions (十二皮部). Collaterals. □. Fifteen collaterals (十五絡). □. Superficial collaterals (浮絡). □. Tertiary collaterals (孫絡)			

**Salviae miltiorrhizae (丹蔘)**

The major function of *Salviae miltiorrhizae* (“Danshen;” 丹蔘) is the activation of blood circulation to dissipate blood stasis. It also offers beneficial effects in the prevention and treatment of cardiovascular diseases [37,38].

**Panax ginseng (人蔘)**

*Panax ginseng* (“Renshen;” 人蔘) has anti-inflammatory, anti-oxidant, and anticancer effects. It also improves psychological and immune function as well as conditions associated with diabetes and is used in the treatment of frailty and aging-related symptoms, such as fatigue and hypertension [39,40].

**Astragalus membranaceus (黃耆)**

*Astragalus membranaceus* (“Huang Qi;” 黃耆) can be used to treat “qi deficiency constitution” [41]. It can also enhance the natural defense mechanisms in the heart, brain, kidney, intestine, liver, and lung [42-44].

**Angelica sinensis (當歸)**

*Angelica sinensis* (“Danggui;” 當歸) used to treat invigorating blood and treating female irregular menstruation and amenorrhea. The *Angelica sinensis* can effectively inhibit inflammation and upregulated growth factor [45].


**12 MAIN MERIDIANS**

In addition to Chinese herbal treatment, acupuncture is another method used to treat AD. The principle of acupuncture is derived from the study of the 12 main meridians and 365 acupuncture points. The 12 main meridians comprise the path of blood and qi that flow through various organs [46]. The meridian system can be further divided into the meridian–collateral system and the tendon–skin system. The meridian–collateral system relates to nutrition and blood (nutrient qi), whereas the tendon–skin system involves the regulation of motor function and body protection (defensive

qi). The meridian–collateral system is composed of 12 principal meridians, eight extra meridians and 15 collaterals. The tendon–skin system includes 12 sinew meridians and 12 cutaneous meridians [47]. The 12 main meridians are distinguished by their yang or yin qi, three levels of energy flow ranging from strong to weak, and direction of flow to either hand or foot [Table 1]. In addition to the 12 meridians, there is the governor vessel (GV) and conception vessel (CV). GV is located on the back, runs from head to hip and has 28 acupuncture points. CV located on the front of the body and has 24 acupuncture points. The six yang meridians have acupuncture points with confluence (i.e., pairing) in CV, and the six yin meridians have acupuncture points with confluence in GV.

**ACUPUNCTURE TREATMENT FOR DEMENTIA**

The bladder (BL) and gallbladder (GB) meridians are very important for the treatment of the brain. The BL meridian has a close relationship with the kidneys and can improve blood supply [Figure 2a and b] [48]. There is a major focus of GB acupuncture points on the head, directing impacting the brain [Figure 2c] [49]. These points are effective in treating dementia [50]. Acupuncture points mark the places where needles can be placed in the body without causing serious harm. Different positions correspond to different disease treatments. The local area comprising an acupuncture point contains a combination of mast cells and blood vessel and nerve networks [Table 2] [51]. Acupuncture points, including Dazhui (GV14), Baihui (GV20), and Shenshu (BL23) are considered to have beneficial effects on dementia. Dazhui is the convergence of all positive meridian points, and Baihui is the convergence of the Tai yang, Shao yang, and Jue yin meridian points. These two acupuncture points are common choices for the treatment of dementia [52]. The term Shenshu means that the kidneys transport cold and moisture to the foot Tai Yang meridian. Dazhui, Baihui, and Shenshu, these three points are related to the kidneys and foot



**Figure 2:** Mapping of the four major meridians directly related to Alzheimer’s disease: bladder, heart, gallbladder, and kidney. The dots indicate acupuncture points along the lines of qi circulation for each meridian. The direction of qi flow is indicated by the arrows. The bladder meridian flows from the head to the back (green line) (a) Kidney, heart, (b) bladder, (c) gallbladder.

**Table 2: Twelve main meridians that connect and regulate the flow of qi through different organs in the body**

Property	Mean	Hand and Foot	Time	Major influence organ	Abbreviation
Tai yin	Yin qi strong	Hand	3-5 o'clock	Lung	LU
		Foot	9-11 o'clock	Spleen	SP
Shao yin	Yin qi weak	Hand	11-13 o'clock	Heart	HT
		Foot	17-19 o'clock	Kidney	KI
Jue yin	Yin qi final step trans to yang qi	Hand	19-21 o'clock	Pericardium	PC
		Foot	1-3 o'clock	Liver	LR
Tai yang	Yang qi strong	Hand	13-15 o'clock	Small intestine	SI
		Foot	15-17 o'clock	Bladder	BL
Shao yang	Yang qi weak	Hand	21-23 o'clock	Triple energizer	TE
		Foot	23-1 o'clock	Gallbladder	GB
Yang ming	Yang qi final step	Hand	5-7 o'clock	Large intestine	LI
		Foot	7-9 o'clock	Stomach	ST

Tai yang meridian [53]. Zusanli (stomach meridian [ST]36) can effectively regulate the physiological functions of the spleen, stomach, and kidney [54]. Baihui and Zusanli were the first choice acupuncture points used treatment dementia [55-57]. Xuehai (spleen meridian [SP]10) and Sishencong (EX-HN 1) are also often used to treat dementia [58].

Alternative acupuncture points to treat dementia include Taixi (Kidney meridian [KI]3) [59], Danzhong (CV17) [60], Zhongwan (CV12) [60], Qihai (CV6), Waiguan (Triple Burner meridian [TE]5), Xuanzhong (GB39), Neiguan (Pericardium meridian [PC]6), Yintang (GV29), Sanyinjiao (SP6), and Dazhong (KI4). The effects of acupuncture on meridians require scientific verification, but analysis of the main meridians is challenging. Fourier analysis can detect the complete quantitative expression of the pressure wave. Wang *et al.* proposed that the detection of harmonic components could

facilitate an understanding of the circulatory system in terms of the different meridians [61,62]. This is a potential method for demonstrating the effects of acupuncture.

### MOXIBUSTION

Methods of treating diseases through acupuncture points include Acupuncture (“zhen,” 針) and moxibustion (“jiu,” 灸). Acupuncture is to insert a needle into acupuncture points to treat diseases, which is an invasive treatment; moxibustion ignites *Artemisia argyi* (“Ai Cao,” 艾草) and through heat generated by cauterization to stimulate the acupuncture points. The function includes adopted to resolve the stagnation, promote circulation in meridian, regulate qi and blood circulation [63]. The moxibustion combine with Typhonium Rhizome (“Bai Fu Zi,” 白附子) was applied to Baihui (BL 20) and the suspended moxibustion was used on Shenting (BL 24) and Dazhui (GV 14). There has a function of improving vascular



dementia and the activity of daily living [64]. In the animal model of AD, moxibustion proved to be helpful for learning and short-term memory capacity [52]. The mechanism of moxibustion treatment of AD is to reduce the content of APP in the frontal cortex and hippocampus [65]. The moxibustion combined with acupuncture is better than the single treatment method to improve the cognitive function and daily activity ability of patients with dementia [66].

## CONCLUSIONS

Given the growing research on the identification of active ingredients from Chinese herbs and their use in the treatment of dementia and cognitive impairment, there are also clinical trials using Chinese herbal medicine and acupuncture to treat AD. If these treatments in effect may be possible to reduce the cost of new drug development and help patients avoid unwanted side effects. This will facilitate the development of future treatments for AD. Furthermore, the combination of herbal medicines with acupuncture may provide additional benefit and could form the basis of a new, synergistic treatment approach for AD.

## Financial support and sponsorship

This research was funded by Buddhist Tzu Chi Medical Foundation, grant number TCMF-SP 109-03.

## Conflicts of interest

Dr. Shinn-Zong Lin, the associate editor of *Tzu Chi Med J*, had no role in the peer review process and the decision to publish this article. The other authors declared that they have no conflicts of interest.

## REFERENCES

- Alzheimer's Association. 2016 Alzheimer's disease facts and figures. *Alzheimers Dement* 2016;12:459-509.
- Long JM, Holtzman DM. Alzheimer Disease: An Update on Pathobiology and Treatment Strategies. *Cell* 2019;179:312-39.
- Chen Y, Zhang W, Wu H, Lao L, Xu J, Xu S. Combination of acupuncture and Chinese herbal formula for elderly adults with mild cognitive impairment: Protocol for a randomized controlled trial. *Trials* 2019;20:117.
- Zhou S, Dong L, He Y, Xiao H. Acupuncture plus herbal medicine for Alzheimer's disease: A systematic review and meta-analysis. *Am J Chin Med* 2017;45:1327-44.
- Hyde AJ, May BH, Dong L, Feng M, Liu S, Guo X, et al. Herbal medicine for management of the behavioural and psychological symptoms of dementia (BPSD): A systematic review and meta-analysis. *J Psychopharmacol* 2017;31:169-83.
- Dawkins E, Small DH. Insights into the physiological function of the beta-amyloid precursor protein: Beyond Alzheimer's disease. *J Neurochem* 2014;129:756-69.
- Sun J, Roy S. The physical approximation of APP and BACE-1: A key event in Alzheimer's disease pathogenesis. *Dev Neurobiol* 2018;78:340-7.
- De Strooper B. Loss-of-function presenilin mutations in Alzheimer disease. *Talking Point on the role of presenilin mutations in Alzheimer disease*. *EMBO Rep* 2007;8:141-6.
- Hardy J. Has the amyloid cascade hypothesis for Alzheimer's disease been proved? *Curr Alzheimer Res* 2006;3:71-3.
- Gendron TF, Petrucelli L. The role of tau in neurodegeneration. *Mol Neurodegener* 2009;4:13.
- Nag S, Yu L, Wilson RS, Chen EY, Bennett DA, Schneider JA. TDP-43 pathology and memory impairment in elders without pathologic diagnoses of AD or FTL. *Neurology* 2017;88:653-60.
- Awada AA. Early and late-onset Alzheimer's disease: What are the differences? *J Neurosci Rural Pract* 2015;6:455-6.
- Yang XM, Li QF, Li DX, Wang YX. Research on developmental characteristic of the theory of ZANG-fu differentiation of syndromes in the period of Jin-Tang Dynasties. *Zhonghua Yi Shi Za Zhi* 2007;37:67-70.
- Chang XL. Synopsis about the hypothesis of "information channel" of channel-collateral system. *Zhen Ci Yan Jiu* 2008;33:356-9.
- Su SB, Jia W, Lu A, Li S. Evidence-based ZHENG: A traditional Chinese medicine syndrome 2013. *Evid Based Complement Alternat Med* 2014;2014:484201.
- Wang HY, Jiang LD, Ma Q, Xu D, Tang SH, Luo ZG. Analysis of clinical characteristics of traditional Chinese and Western medicine in professor Jiang Liangduo's theory of "sanjiao meridian stasis". *Zhongguo Zhong Yao Za Zhi* 2017;42:4875-81.
- Wax WA, White KR. A short history of a long tradition. Qi: The basis of Chinese medicine. *Am J Nurs* 2000;100:42-3.
- Culliton BJ. NIH says "yes" to acupuncture. *Nat Med* 1997;3:1307.
- McAdams-DeMarco MA, Bae S, Chu N, Gross AL, Brown CH, Oh E, et al. Dementia and Alzheimer's disease among older kidney transplant recipients. *J Am Soc Nephrol* 2017;28:1575-83.
- Wang X, Wang Y, Yu S, Ren L. Effect of electroacupuncture on levels of  $\beta$ -amyloid and neprilysin proteins in the cerebral cortex of Alzheimer's disease mice based on "mutual assistance of kidney and brain" theory. *Zhen Ci Yan Jiu* 2018;43:20-4.
- Ong WY, Wu YJ, Farooqui T, Farooqui AA. Qi fu yin-a ming dynasty prescription for the treatment of dementia. *Mol Neurobiol* 2018;55:7389-400.
- SZ. *The Divine Farmer's MateriaMedica: A Translation of the Shen Nong Ben Cao Jing*. Boulder, CO: Blue Poppy Press; 1998.
- Chen R. *Medical thoughts in Ben Cao Gang Mu (Compendium of materia medica)* (Chi). *Zhonghua Yi Shi Za Zhi*, 1987;17:42-5.
- Cho JH, Jung JY, Lee BJ, Lee K, Park JW, Bu Y. *Epimedii herba: A promising herbal medicine for neuroplasticity*. *Phytother Res* 2017;31:838-48.
- Ulbricht CE, Natural Standard Research Collaboration. An evidence-based systematic review of yin yang huo (*Epimedium spp.*) by the natural standard research collaboration. *J Diet Suppl* 2016;13:136-64.
- Shin IJ, Son SU, Park H, Kim Y, Park SH, Swanberg K, et al. Preclinical evidence of rapid-onset antidepressant-like effect in *Radix Polygalae* extract. *PLoS One* 2014;9:e88617.
- Liu C, Zhang A, Yan GL, Shi H, Sun H, Han Y, et al. High-throughput ultra high performance liquid chromatography coupled to quadrupole time-of-flight mass spectrometry method for the rapid analysis and characterization of multiple constituents of *Radix Polygalae*. *J Sep Sci* 2017;40:663-70.
- Yi YQ, Fang R, Ge JW, Cheng SW, Wang GZ, Liu L. Analysis on medication rules for treatment of dementia by ancient physicians based on data mining methods. *Zhongguo Zhong Yao Za Zhi* 2018;43:3376-81.
- Soon YY, Tan BK. Evaluation of the hypoglycemic and anti-oxidant activities of *Morinda officinalis* in streptozotocin-induced diabetic rats. *Singapore Med J* 2002;43:077-85.
- Zhang FH, Wang ZM, Liu YT, Huang JS, Liang S, Wu HH, et al. Bioactivities of serotonin transporter mediate antidepressant effects of *Acori tatarinowii* Schott. *J Ethnopharmacol* 2019;241:111967.
- Song Z, Yin F, Xiang B, Lan B, Cheng S. Systems pharmacological approach to investigate the mechanism of *Acori Tatarinowii Rhizoma* for Alzheimer's disease. *Evid Based Complement Alternat Med* 2018;2018:5194016.
- Wu QD, Yuan DJ, Wang QW, Wu XR. Effects of volatile oil of *Rhizoma Acori Tatarinowii* on morphology and cell viability in cultured cardiac myocytes. *Zhong Yao Cai* 2009;32:242-5.
- Zhang RX, Li MX, Jia ZP. *Rehmannia glutinosa*: Review of botany, chemistry and pharmacology. *J Ethnopharmacol* 2008;117:199-214.

34. Lee B, Shim I, Lee H, Hahm DH. Rehmannia glutinosa ameliorates scopolamine-induced learning and memory impairment in rats. *J Microbiol Biotechnol* 2011;21:874-83.
35. Feng ZY, Cui Y, Ding G, Zhang B. Experimental research about the effect of Rehmannia glutinosa on the drug effect fingerprints of the rabbit's blood serum of Yin deficiency and heat flourishing syndrome. *Zhong Yao Cai* 2007;30:1545-8.
36. Yang R, Yuan BC, Ma YS, Zhou S, Liu Y. The anti-inflammatory activity of licorice, a widely used Chinese herb. *Pharm Biol* 2017;55:5-18.
37. Naveed M, Wenhua L, Gang W, Mohammad IS, Abbas M, Liao X, et al. A novel ventricular restraint device (ASD) repetitively deliver *Salvia miltiorrhiza* to epicardium have good curative effects in heart failure management. *Biomed Pharmacother* 2017;95:701-10.
38. Zhang L, Gan W, An G. Influence of Tanshinone IIA on heat shock protein 70, Bcl-2 and Bax expression in rats with spinal ischemia/reperfusion injury. *Neural Regen Res* 2012;7:2882-8.
39. Shergis JL, Di YM, Zhang AL, Vlahos R, Helliwell R, Ye JM, et al. Therapeutic potential of *Panax ginseng* and ginsenosides in the treatment of chronic obstructive pulmonary disease. *Complement Ther Med* 2014;22:944-53.
40. Ogawa-Ochiai K, Kawasaki K. *Panax ginseng* for frailty-related disorders: A review. *Front Nutr* 2018;5:140.
41. Adesso S, Russo R, Quaroni A, Autore G, Marzocco S. *Astragalus membranaceus* extract attenuates inflammation and oxidative stress in intestinal epithelial cells via NF- $\kappa$ B activation and Nrf2 response. *Int J Mol Sci* 2018;19:800.
42. Huang YC, Tsay HJ, Lu MK, Lin CH, Yeh CW, Liu HK, et al. *Astragalus membranaceus*-polysaccharides ameliorates obesity, hepatic steatosis, neuroinflammation and cognition impairment without affecting amyloid deposition in metabolically stressed APP<sup>swe</sup>/PS1<sup>dE9</sup> Mice. *Int J Mol Sci* 2017;18:2746.
43. Cho WC, Leung KN. *In vitro* and *In vivo* immunomodulating and immunorestorative effects of *Astragalus membranaceus*. *J Ethnopharmacol* 2007;113:132-41.
44. Shahzad M, Shabbir A, Wojcikowski K, Wohlmuth H, Gobe GC. The antioxidant effects of radix *astragali* (*Astragalus membranaceus* and Related Species) in protecting tissues from injury and disease. *Curr Drug Targets* 2016;17:1331-40.
45. Duan MH, Wang LN, Jiang YH, Pei YY, Guan DD, Qiu ZD. *Angelica sinensis* reduced A $\beta$ -induced memory impairment in rats. *J Drug Target* 2016;24:340-7.
46. Li SY, Zhao JS. Evolution of the connotation of meridians viewed from the changes in the traveling routes: Considerations induced from the "eleven meridians" to the "twelve regular meridians". *Zhen Ci Yan Jiu* 2009;34:132-5.
47. Liu N. Hypothesis of "sinew-meridian system". *Zhongguo Zhen Jiu* 2017;37:79-83.
48. Yao F, Zhao Y, Jiang S, Fang M. The theoretical basis for chronic fatigue syndrome from bladder meridian of foot-taiyang. *Zhongguo Zhen Jiu* 2015;35:295-8.
49. Lin XM, Yao X, Di Z. Acupuncture at "Siguan" combined with Gallbladder Meridian acupoints for migraine: A randomized controlled trial. *Zhongguo Zhen Jiu* 2014;34:947-50.
50. Chen XJ, Chen LF, Chen Q, Fang Z. Effect of three-line puncture on the Governor Vessel and Bladder Meridian on head on cognitive function of vascular dementia patients. *Zhongguo Zhen Jiu* 2012;32:289-92.
51. Zhao P, Wang Y, Gu F, Li C, Wei Y, Wang G, et al. Comparison of the effects of the intervention with electric thermal bian stone and air suction cup on blood perfusion at meridian points. *Zhongguo Zhen Jiu* 2018;38:159-64.
52. Zhu CF, Zhang LD, Song XG, Yang J, Pan HP, He CG, et al. Moxibustion improves learning-memory ability by promoting cellular autophagy and regulating autophagy-related proteins in hippocampus and cerebral cortex in APP/PS1 transgenic Alzheimer's disease mice. *Zhen Ci Yan Jiu* 2019;44:235-41.
53. Ye Y, Zhu W, Wang XR, Yang JW, Xiao LY, Liu Y, et al. Mechanisms of acupuncture on vascular dementia-A review of animal studies. *Neurochem Int* 2017;107:204-10.
54. Liu J, Liu J, Wang G, Liu G, Zhou H, Fan Y, et al. Electroacupuncture at Guanyuan (CV 4), Zusanli (ST 36) and Baihui (DU 20) regulate the aging-related changes in gene expression profile of the hippocampus in sub-acutely aging rats. *PLoS One* 2018;13:e0191623.
55. Zhou J, Peng W, Xu M, Li W, Liu Z. The effectiveness and safety of acupuncture for patients with Alzheimer disease: A systematic review and meta-analysis of randomized controlled trials. *Medicine (Baltimore)* 2015;94:e933.
56. Tian GH, Sun K, Huang P, Zhou CM, Yao HJ, Huo ZJ, et al. Long-term stimulation with electroacupuncture at du20 and st36 rescues hippocampal neuron through attenuating cerebral blood flow in spontaneously hypertensive rats. *Evid Based Complement Alternat Med* 2013;2013:482947.
57. Kan BH, Yu JC, Zhao L, Zhao J, Li Z, Suo YR, et al. Acupuncture improves dendritic structure and spatial learning and memory ability of Alzheimer's disease mice. *Neural Regen Res* 2018;13:1390-5.
58. Yu J, Zhang X, Liu C, Meng Y, Han J. Effect of acupuncture treatment on vascular dementia. *Neurol Res* 2006;28:97-103.
59. Zhu B, Wang Y, Zhang G, Ouyang H, Zhang J, Zheng Y, et al. Acupuncture at KI3 in healthy volunteers induces specific cortical functional activity: An fMRI study. *BMC Complement Altern Med* 2015;15:361.
60. Yu J, Liu C, Zhang X, Han J. Acupuncture improved cognitive impairment caused by multi-infarct dementia in rats. *Physiol Behav* 2005; 86:434-41.
61. Chang CW, Wang WK. Reliability assessment for pulse wave measurement using artificial pulse generator. *J Med Eng Technol* 2015; 39:177-84.
62. Chang CW, Wang SH, Jan MY, Wang WK. Effect of black tea consumption on radial blood pulse spectrum and cognitive health. *Complement Ther Med* 2017;31:1-7.
63. Litscher G. Bioengineering assessment of acupuncture, part 8: Innovative moxibustion. *Crit Rev Biomed Eng* 2010;38:117-26.
64. Chen H, Wang P, Yang J, Liu G. Impacts of moxibustion on vascular dementia and neuropeptide substance content in cerebral spinal fluid. *Zhongguo Zhen Jiu* 2011;31:19-22.
65. Wang JJ, Wu SB, Cheng HL, Yang C, Chen Y, Yang J. Effect of moxibustion on expression of RAGE and LRP-1 and neuronal ultrastructure of frontal cortex and hippocampus in vascular dementia rats. *Zhen Ci Yan Jiu* 2020;45:33-9.
66. Wang S, Ye G, Xu C, Jia A, Ru Y, Guan S, et al. Flipping moxibustion of Hu medicine at acupoints in governor vessel combined with acupuncture for vascular dementia. *Zhongguo Zhen Jiu* 2018;38:919-24.